



19(6)

XVII Iberian Symposium on
Marine Biology Studies
(SIEBM)

11th-14th September, 2012.
Donostia-San Sebastián (Spain)



Ángel Borja
(Editor)

Borja, A. (Editor), 2012. XVII Iberian Symposium on Marine Biology Studies. Revista de Investigación Marina, AZTI-Tecnalia, 19(6): 108-631.

La serie 'Revista de Investigación Marina', editada por la Unidad de Investigación Marina de Tecnalia, cuenta con el siguiente Comité Editorial:

Editor: Dr. Ángel Borja

Adjunta al Editor: Dña. Mercedes Fernández Monge e Irantzu Zubiaur (coordinación de las publicaciones)

Comité Editorial: Dr. Lorenzo Motos
Dr. Adolfo Uriarte
Dr. Michael Collins
Dr. Javier Franco
D. Julien Mader
Dña. Marina Santurtun
D. Victoriano Valencia
Dr. Xabier Irigoien
Dra. Arantza Murillas
Dr. Josu Santiago

La 'Revista de Investigación Marina' de Tecnalia edita y publica investigaciones y datos originales resultado de la Unidad de Investigación Marina de Tecnalia. Las propuestas de publicación deben ser enviadas al siguiente correo electrónico aborja@azti.es. Un comité de selección revisará las propuestas y sugerirá los cambios pertinentes antes de su aceptación definitiva.



Edición: 1.ª Septiembre 2012

© AZTI-Tecnalia

ISSN: 1988-818X

Unidad de Investigación Marina

Internet: www.azti.es

Edita: Unidad de Investigación Marina de Tecnalia

Herrera Kaia, Portualdea

20010 Pasaia

Foto portada: Iñigo Onandia (vista aérea del barco *Leporre Anaiak*, de Bermeo) © AZTI-Tecnalia.

© AZTI-Tecnalia 2012. Distribución gratuita en formato PDF a través de la web: www.azti.es/RIM

Virgin biomass, fishing potential and valorisation of *Plesionika edwardsii* (Crustacea: Decapoda: Caridea: Pandalidae) in the Cape Verde Islands - Preliminary results

J.A. González^{a,b}, O. Tariche^c, N. Almeida^d, C. Monteiro^c, S. Jiménez^a, A. Martins^c, J.I. Santana^a y J.G. Pajuelo^{a,d}

These studies are part of the PROACTIVA 1-2 (2009-2012) and MARPROF-CV projects (2010-2013), in the framework of the Canary Islands Government and UE PCT MAC 2007-2013 programmes respectively. Research has mainly focused on the stock assessment of the striped soldier shrimp, *Plesionika edwardsii* (Brandt, 1851), because it has shown moderate to high levels of fishing yield and abundances compatible with the development of a new sustainable fishery in the Cape Verde Islands (González & Tariche, eds., 2009). Several actions have been done in order to implement an ecosystem approach. Other aims have prosecuted to contribute to the professional training through technology transfer actions, and to promote and disseminate new fish products through public presentations and valorisation gastronomic events reinforced with an informative exhibition.

Cruises took place onboard the R/V “Profesor Ignacio Lozano”. Of the four cruises scheduled, three 15-day cruises have been conducted to date: Camarão-0 (around São Vicente and Santa Luzia islands, April 2010) within the PROACTIVA 1 project, Camarão-1 (Santiago island, November 2011) and Camarão-2 (Boa Vista island, Mars 2012) within the MARPROF-CV project. A standardized fishing gear was used, so-called multiple semifloating shrimp traps (MSFST), each fishing line with 40-65 traps operating around 2.4 m above the seafloor (González *et al.* 1992). The most abundant, coastal pelagic fish in the Cape Verde waters, the so-called “cavala preta” *Decapterus macarellus* (Cuvier, 1833) (Carangidae), was used as unique bait of the traps.

An echo-sounding bathymetric survey was done followed by a prospection around the mentioned islands, and finally a depletion model approach was applied. Temperature and salinity profiles have been obtained by the use of CTD during fishing operations and these data were further related with the relative abundance indices of the target species.

Initial biomass was estimated from depletion experiments at different target species’ abundance stations by applying the Leslie & Davis (1939) method, adapted and modified by Ricker

(1975). Density by area was calculated assuming two different areas of attraction of the fishing gear. Each trap was attached to the main bottom line every 15 m (González *et al.* 1992), and the maximum attraction of this bait was established to be 100 or 150 m, according to the experience gained from the local fishery for lobsters. Each fishing operation was classified according to its yield (CPUE in g/trap/night). The potential fishing planar area was estimated between the isobathymetric lines between 90 and 220 m. Total biomasses (Bt) were calculated from areas (km²) and mean minimum/maximum densities (kg/km²).

Maximum sustainable yield (MSY) was estimated from Bt using the Beddington and Cooke (1983) model by entering natural mortality (0.6), growth rate (0.53 year⁻¹) and recruitment age (1.32 year) parameters for the target species published for the Canary Islands (Santana *et al.* 1997, Quiles, 2005). These parameters estimate an exploitation rate (β) of 0.262. Based on a quarterly work strategy, the life cycle of this target species in the Cape Verdes is being studied in order to check the stock estimations.

The MSY estimated to date were 30.5 tons/year for the stock of São Vicente, Santa Luzia, Ilheus and São Nicolau islands, 10 tons/year for the stock of Santiago island, and 139 tons/year for the stock of Boa Vista and Maio islands. In a near future, the stock of the Sal island will be equally assessed, meanwhile the small stocks of Santo Antão, Fogo and Brava islands will be estimated by interpolation. More than 200 tons per year is expected to be the total MSY for the striped soldier shrimp around the islands of the Cape Verde archipelago, occupying a total area of more than 1,900 km² of new fishing grounds at between 90 and 220 m of depth.

In comparison with the metallic bottom traps, which are traditional and intensively used in the Canary Islands, the innovative fishing gear MSFST seems to be more selective for pandalid shrimps, minimizing the gear impact on the seafloor as well as the by-catch by reducing the discards.

Depletion methods are based on the assumption of a closed system. The straightforward decline of CPUEs obtained during the depletion experiments seems to confirm that *P. edwardsii* is a low mobility species, making this assumption valid at least during short-time periods. Because of the bathymetry profile of these islands, the depth range is from very close (few nautical miles in Santiago to far away (10-12 miles in Boa Vista) to the coastline. Additionally, all specimens caught have been identified, counted and weighted at species level, ecological studies, vertical distribution and basic biological parameters of several by-catch species involved in the future fishery are being studied.

^a Grupo de Biología Pesquera, Instituto Canario de Ciencias Marinas (ICCM-ACIISI), Gobierno de Canarias (solea@iccm.recanaria.es).

^b Grupo de Ecología Marina Aplicada y Pesquerías, Universidad de Las Palmas de Gran Canaria (ULPGC).

^c Instituto Nacional de Desenvolvimento das Pescas (INDP) de Cabo Verde.

^d Departamento de Engenharia e Ciências do Mar (DECM), Universidade de Cabo Verde (Uni-CV).

In parallel, according to the communication plan scheduled (González & Tariche, eds., 2009), local biologists, technologists, technicians and fishermen have been trained in the fishing operations onboard both the research vessel and the local fishing boat “Sinagoga”, in the biological sampling at laboratory, and by means a theoretical and practical course for hand-making traps and the whole fishing gear for 30 people. After every campaign, main results were presented and discussed with the local stakeholders, followed by a culinary laboratory and a gastronomic tasting based on target and by-catch species, which were also valorised with nutritional biochemical analyses. All these actions were reinforced with an informative exhibition composed of four panels of large format and several explanation posters and displays (roll-ups), as well as by means radio and TV interviews.

When biological and assessment studies were concluded, the management options using the best methods and data available should be analyzed, especially prior to fishing activity. Fishing effort should be controlled on the basis of quotas, number of fishing vessels and a precautionary approach in order to ensure that catching is commensurate with sustainable levels of exploitation. The MSY estimates suggest that this new Capeverdean fishery should be carried out by several specialized medium-sized fishing vessels, fitting well with the current artisanal fleets of the Macaronesian small-scale fisheries.

During the last decades a combination of shrimp trawling and industrial trapping activity has threatened over-exploitation in the Mediterranean fisheries targeting on *P. edwardsii*. Currently the shrimp collapse has conducted to the decline of these fisheries. Can the Cape Verde regulatory bodies and all the stakeholders involved learn the lessons this teaches us about this resource management?

References

- Beddington, J.R. & J.G. Cooke (1983) The potential yield of fish stocks. *FAO Fish. Tech. Pap.*, 242: 47 pp.
- González, J.A. & O. Tariche (eds.) (2009) Una mirada sobre la biodiversidad marina y bases para su gestión sostenible. Recursos pesqueros potenciales de profundidad de Cabo Verde. Dirección General de Relaciones con África del Gobierno de Canarias. Las Palmas de Gran Canaria: 176 pp.
- González, J.A., J. Carrillo, J.I. Santana, P. Martínez Baño & F. Vizuete (1992) La pesquería de Quisquilla, *Plesionika edwardsii* (Brandt, 1851), con tren de nasas en el Levante español. Ensayos a pequeña escala en Canarias. *Inf. Téc. Sci. Mar.*, 170: 31 pp.
- Leslie, P.H. & D.H.S. Davis (1939) An attempt to determine the absolute number of rats on a given area. *J. An. Ecol.*, 8: 94-113.
- Quiles, J.A. (2005) Biología, evaluación y plan piloto de pesca del stock de camarón soldado *Plesionika edwardsii* (Crustacea, Decapoda, Pandalidae) de Gran Canaria. Tesis doctoral. Universidad de Las Palmas de Gran Canaria, Facultad de Ciencias del Mar.
- Ricker, W.E. (1975) Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Res. Bd Can.*, 191: 382 pp.
- Santana, J.I., J.A. González, I.J. Lozano & V.M. Tuset (1997) Life history of *Plesionika edwardsii* (Crustacea, Decapoda, Pandalidae) around the Canary Islands (Eastern Central Atlantic). *S. Afr. J. mar. Sci.*, 18: 39-48.



www.azti.es

azti
tecnalia

Txatxaramendi ugartea z/g
48395 Sukarrieta (Bizkaia)
Tel.: +34 94 657 40 00
Fax: +34 94 657 25 55

Herrera Kaia, Portualdea z/g
20110 Pasaia (Gipuzkoa)

Parque Tecnológico de Bizkaia
Astondo bidea. Edificio 609.
48160 Derio (Bizkaia)